

## REPORT DOCUMENTATION PAGE

0077

0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including gathering and maintaining the data needed, and completing and reviewing the collection of information, including suggestions for reducing this burden, to Washington Headquarters, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

ing data sources.  
her aspect of this  
ts, 1215 Jefferson

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT TYPE AND DATES COVERED Final 15 Feb 97 To 14 Aug 97	
4. TITLE AND SUBTITLE 1997 COUPLED CLUSTER AND ELECTRON CORRELATION WORKSHOP "FIFTY YEARS OF THE CORRELATION PROBLEM"				5. FUNDING NUMBERS F49620-97-1-0091 2303/FS 61102F	
6. AUTHOR(S) Dr Rodney J. Bartlett					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Dept of Chemistry & Physics University of Florida P.O. Box 118435 Gainesville FL 32611-8435				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NL 110 Duncan Ave Room B115 Bolling AFB DC 20332-8050  Dr Michael R. Berman				10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION AVAILABILITY STATEMENT  Approved for public release; distribution unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The Coupled Cluster Theory Electron Correlation Workshop "Fifty Years of the Correlation Problem" was held at Cedar Key, Florida, from June 15-19, 1997, to recognize the essential developments for one of the dominant topics in the quantum theory of atoms, molecules, and solids. The instantaneous Coloumbic interactions among electrons that correlate their motion (the electron correlation problem) has been the focal point of ab initio quantum chemistry and physics for many years. Only with the proper inclusion of electron correlation in approximate solutions of the Schroedinger (or Dirac-Fock) equation is it possible to provide predictive accuracy for most properties of atoms and molecules. Such quantities include energetics (involving multiplets, dissociation pathways, and activation barriers), excited states and first- and second-order properties (like moments, field, gradients, polarizabilities, and magnetic susceptibilities), and vibrational, electronic, EPR and NMR spectra, among others.					
14. SUBJECT TERMS				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT (U)	18. SECURITY CLASSIFICATION OF THIS PAGE (U)	19. SECURITY CLASSIFICATION OF ABSTRACT (U)	20. LIMITATION OF ABSTRACT (UL)		

19980116 100

## **FINAL REPORT—AFOSR F49620-97-1-0091**

### **THE COUPLED CLUSTER THEORY ELECTRON CORRELATION WORKSHOP "FIFTY YEARS OF THE CORRELATION PROBLEM"**

The Coupled Cluster Theory Electron Correlation Workshop "Fifty Years of the Correlation Problem" was held at Cedar Key, Florida, from June 15-19, 1997, to recognize the essential developments for one of the dominant topics in the quantum theory of atoms, molecules, and solids. The instantaneous Coloumbic interactions among electrons that correlate their motion (the electron correlation problem) has been the focal point of *ab initio* quantum chemistry and physics for many years. Only with the proper inclusion of electron correlation in approximate solutions of the Schroedinger (or Dirac-Fock) equation is it possible to provide predictive accuracy for most properties of atoms and molecules. Such quantities include energetics (Involving multiplets, dissociation pathways, and activation barriers), excited states and first- and second-order properties (like moments, field, gradients, polarizabilities, and magnetic susceptibilities), and vibrational, electronic, EPR and NMR spectra, among others.

In extended systems like polymers and solids, correlation effects are critical to properties like cohesive energies, energy bands, and particularly band gaps; other optical properties like exciton spectra or non-linear optical behavior; and collective phenomena as in conductivity and superconductivity. Without adequate inclusion of electron correlation, theory cannot be predictive.

About 50 years after the correlation problem was clearly identified, we were in a position to report on the remarkable progress that has been made and to point the way to the future. These accomplishments have been achieved by a combination of critical formal and methodological developments in the theory such as coupled-cluster theory, many-body perturbation theory, propagator methods, and of configuration interaction methods coupled to the dramatic progress in computational power. Together, today, accurate applications to molecular many-electron systems are possible. This meeting addressed developing a similar capability for the reliable introduction of electron correlation to extended systems.

To achieve some historical perspective, several of the pioneer investigators who have made contributions to the correlation problem spoke at the meeting. This, in fact, enabled us to have representatives of the four scientific generations of investigators who have worked on the correlation problem.

The meeting was attended by 88 scientists from over 30 different countries. Thirty-seven invited presentations were made, augmented by 37 poster presentations. In a panel session, we addressed the timely topic of "With Density Functional Theory, Is there Any Future for *ab initio* Correlated Methods?" which generated much discussion.

The Proceedings, which consists of 25 papers, will be published in *Molecular Physics*. Copies of the Proceedings will be sent to AFOSR as instructed when they become available.